

Few-atom approach to many-body physics


G.J. Conduit and P.O. Bugnion

POB, J. Lofthouse & GJC, Phys. Rev. Lett. **111**, 045301 (2013)

POB & GJC, Phys. Rev. A **87**, 060502(R) (2013)

POB & GJC, Phys. Rev. A **88**, 013601 (2013)

Experimental setup



$|F = 1/2, m_F = 1/2\rangle$



Up spin electron


$|F = 1/2, m_F = -1/2\rangle$



Down spin electron


$$\hat{H} = -\frac{\nabla^2}{2} + g n_{\uparrow}(\vec{r}) n_{\downarrow}(\vec{r})$$

Two distinguishable fermions




G. Zürn *et al.* PRL 108 075303 (2012)


Two distinguishable fermions




Two distinguishable fermions



Two distinguishable fermions




Energy of states




Weak
repulsion

Strong
repulsion




Polaron state




J.B. McGuire, J. Math. Phys. **6**, 432 (1965)

Polaron state




P.O. Bugnion & G.J. Conduit Phys. Rev. A **87**, 060502(R) (2013)

Polaron state



A. Wenz *et al.*, arXiv:1307.3443

Polaron state



Strongly correlated physics

Repulsive interactions → Ferromagnetic correlations

Attractive interactions → BCS pairing / FFLO

Double well → Exchange interactions


Zero dimensional → Hund's rules

Perturbed trap → Rabi / Josephson oscillations


Stoner Hamiltonian

$$\hat{H} = -\frac{\nabla^2}{2} + g n_{\uparrow}(\vec{r}) n_{\downarrow}(\vec{r}) + V(\vec{r})$$


Density profiles




Density profiles




Polaron state




Polaron state




Polaron state





Polaron state




Polaron state




Loss region




Tunneling probability




Why probability of $\frac{1}{2}$?



Why probability of $\frac{1}{2}$?




Why probability of $\frac{1}{2}$?




Two-atom scattering

Repulsive




Two-atom scattering

Repulsive




Attractive




Two-atom scattering


Repulsive



Attractive




Repulsive




Two-atom scattering


Repulsive





Attractive





Repulsive





Three-atom bound state





Three-atom bound state





Three-atom bound state





Three-atom bound state




Three-atom bound state





Three-atom bound state





Three-atom bound state




Three-atom bound state





Three-atom bound state




Tunneling probability



Band crossings




Band crossings




BCS Hamiltonian

$$\hat{H} = -\frac{\nabla^2}{2} + g c_{\vec{p}\uparrow}^\dagger c_{-\vec{p}\downarrow}^\dagger c_{-\vec{k}\downarrow} c_{\vec{k}\uparrow} + V(\vec{r})$$


Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing





Inhomogeneous pairing




Inhomogeneous pairing



Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing


(2,1) (3,2) (4,3)




(3,1) (4,2)




(4,1)




$(2,1)^s$ (5,1)



$(2,1)^d$ (5,2)




$(2,1)^g$




Inhomogeneous pairing


Weak interactions



Strong interactions




Inhomogeneous pairing




P.O. Bugnion & GJC, Phys. Rev. A 88, 013601 (2013)

Inhomogeneous pairing



P.O. Bugnion & GJC, Phys. Rev. A 88, 013601 (2013)

Inhomogeneous pairing




Summary

A few-fermion system provides insight into many-body physics

Discretization of energy levels means that losses occur in narrow range of interaction strengths


Observation of Fermi surface and magnetic correlations

Other phenomena



Hund's rules


P.O. Bugnion, J.A. Lofthouse & GJC




Exchange interactions

P.O. Bugnion & GJC
Phys. Rev. A **88**, 013601 (2013)

Finding the missing probability




Finding the missing probability





Inhomogeneous pairing





Inhomogeneous pairing





Inhomogeneous pairing





Inhomogeneous pairing




Inhomogeneous pairing




Inhomogeneous pairing



Inhomogeneous pairing



Inhomogeneous pairing



Inhomogeneous pairing

