

# Few-atom approach to many-body physics


G.J. Conduit and P.O. Bugnion

POB, J. Lofthouse & GJC, Phys. Rev. Lett. **111**, 045301 (2013)

POB & GJC, Phys. Rev. A **87**, 060502(R) (2013)


POB & GJC, Phys. Rev. A **88**, 013601 (2013)

# Two distinguishable fermions




G. Zürn *et al.* PRL 108 075303 (2012)


# Two distinguishable fermions




# Two distinguishable fermions



# Two distinguishable fermions




# Energy of states





Weak  
repulsion

Strong  
repulsion




# Polaron state




$\infty$  atom

2 atom



J.B. McGuire, J. Math. Phys. **6**, 432 (1965)


# Polaron state



P.O. Bugnion & G.J. Conduit Phys. Rev. A **87**, 060502(R) (2013)




# Polaron state




A. Wenz *et al.*, arXiv:1307.3443


# Polaron state




# Polaron state




# Polaron state




# Polaron state




# Polaron state



# Polaron state




# Three-atom bound state







# Three-atom bound state




# Three-atom bound state




# Three-atom bound state




# Three-atom bound state




# Three-atom bound state




# Three-atom bound state



# Three-atom bound state





# Three-atom bound state







# Loss region



# Tunneling probability




# Other phenomena



Inhomogeneous pairing

P.O. Bugnion, J.A. Lofthouse & GJC  
Phys. Rev. Lett. in press



Exchange interactions

P.O. Bugnion & GJC  
Phys. Rev. A **88**, 013601 (2013)

# Summary

A few-fermion system provides insight into many-body physics

Discretization of energy levels inhibits losses, allowing observation of magnetic correlations